

References

1. National Center for Health Statistics. Health, United States, 2008 with special feature on the health of young adults. *Hyattsville, MD: National Center for Health Statistics 2009.*
2. Merikangas KR, He J, Burstein M, Swanson SA, Avenevoli S, Cui L, et al. Lifetime prevalence of mental disorders in U.S. adolescents: Results from the National Comorbidity Study-Adolescent Supplement (NCS-A). *Journal of the American Academy of Child and Adolescent Psychiatry.* 2010; 49(10): 980-989.
3. Corscadden L, Taylor A, Sebold A, Maddocks E, Pearson C, Harvey J, et al. *Obesity in Canada: A joint report from the Public Health Agency of Canada and the Canadian Institute for Health Information.* Public Health Agency of Canada. 2011.
4. Public Health Agency of Canada. *What is depression?* <http://www.phac-aspc.gc.ca/cd-mc/mi-mm/depression-eng.php> (accessed 28 February 2014).
5. World Health Organization . *Adolescent development.* http://www.who.int/maternal_child_adolescent/topics/adolescence/dev/en/ (accessed 28 February 2014).
6. Anderson SE, Murray DM, Johnson CC, Elder JP, Lytle L, Jobe JB, et al.. Obesity and depressed mood associations differ by race/ethnicity in adolescent girls. *International Journal of Pediatric Obesity* 2011; 6(1): 69-78.
7. Boutelle KN, Hannan P, Fulkerson JA, Crow SJ, Stice E. Obesity as a prospective predictor of depression in adolescent females. *Health Psychology* 2010; 29(3): 293-298.
8. Daniels J. Weight and weight concerns: Are they associated with reported depressive symptoms in adolescents? *Journal of Pediatric Health Care* 2005; (19): 33-41.
9. Goodman E, Whitaker RC. A prospective study of the role of depression in the development and persistence of adolescent obesity. *Pediatrics* 2002; 109(3).
10. Halfon N, Larson K, Slusser W. Associations between obesity and comorbid mental health, developmental, and physical health conditions in a nationally representative sample of US children aged 10 to 17. *Academic Pediatrics* 2013; 13(1): 6-13.
11. Kubzansky LD, Gilthorpe MS, Goodman E. A prospective study of psychological distress and weight status in adolescents/young adults. *Annals of Behavioral Medicine* 2012; 43(2): 219-228.
12. Merikangas AK, Mendola P, Pastor PN, Reuben CA, Cleary SD. The association between major depressive disorder and obesity in US adolescents: Results from the 2001-2004 National Health and Nutrition Examination Survey. *Journal of Behavioral Medicine* 2012; (35): 149-154.

13. Mond J, van den Berg P, Boutelle K, Hannan P, Neumark-Sztainer D. Obesity, body dissatisfaction, and emotional well-being in early and late adolescence: Findings from the Project EAT Study. *Journal of Adolescent Health* 2011; (48): 373-378.
14. Richardson LP, Garrison MM, Drangsholt M, Mancl L, LeResche L. Associations between depressive symptoms and obesity during puberty. *General Hospital Psychiatry* 2006; (28): 313-320.
15. Rieck T, Jackson A, Martin SB, Petrie T, Greenleaf C. Health-related fitness, body mass index, and risk of depression among adolescents. *Official Journal of the American College of Sports Medicine* 2013; (): 1083-1088.
16. Roberts RE, Duong HT. Perceived weight, not obesity, increases risk for major depression among adolescents. *Journal of Psychiatric Research* 2013; (47): 1110-1117.
17. Stice E, Presnell K, Shaw H. Psychological and behavioral risk factors for obesity onset in adolescent girls: A prospective study. *Journal of Consulting and Clinical Psychology* 2005; 73(2): 195-202.
18. Swallen KC, Reither EN, Haas SA, Meier AM . Overweight, obesity and health-related quality of life among adolescents: the National Longitudinal Study of Adolescent Health. *Pediatrics* 2005; 115(2): 340-347.
19. Public Health Agency of Canada. *Obesity in Canada – Snapshot*. 2012.
20. World Health Organization. *BMI-for-age (5-19 years)*.
http://www.who.int/growthref/who2007_bmi_for_age/en/ (accessed 5 March 2014).